

Anti-Abscisic acid 8'-hydroxylase 2, C-terminal antibody

Catalog: PHY1832A

Product Information

Description:	Rabbit polyclonal antibody
Background:	CYP707A2 is a protein with ABA 8'-hydroxylase activity, involved in ABA catabolism. It appears to play a major role in the rapid decrease in ABA levels during early seed imbibition.
Synonyms:	CYP707A2, CYTOCHROME P450, FAMILY 707, SUBFAMILY A, POLYPEPTIDE 2
Immunogen:	KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> CYP707A2 (AT2G29090).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4). "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	55 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Brassica rapa</i> , <i>Brassica napus</i> , and 80-99% homologues with the sequence in

Research Use Only

Gossypium raimondii, *Spinacia oleracea*, *Glycine max*, *Populus trichocarpa*, *Cucumis sativus*.

For more species homologues information, please contact tech support at tech@phytoab.com.