

## Anti-Serine/threonine-protein kinase STN7, chloroplastic antibody

Catalog: PHY2832A

## **Product Information**

**Description:** Rabbit polyclonal antibody

Background: STN7 is required for state transitions, phosphorylation of the major antenna

complex (LHCII) between PSII and PSI, and light adaptation.

**Synonyms:** STN7, STT7 HOMOLOG STN7

Immunogen: KLH-conjugated synthetic peptide (20 aa from Central section) derived from

Arabidopsis thaliana STN7 (AT1G68830).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 63 / 54 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in *Vitis* vinifera, Glycine max, Gossypium raimondii, Zea mays, Cucumis



sativus, Populus trichocarpa, Spinacia oleracea, Nicotiana tabacum, Medicago truncatula, Brassica napus, Brassica rapa, Setaria viridis, Sorghum bicolor, Panicum virgatum, Hordeum vulgare, Triticum aestivum, Oryza sativa, Solanum tuberosum, Solanum lycopersicum, and 80-99% homologues with the sequence in Chlamydomonas reinhardtii.

For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Thy-WT: thylakoid membrane proteins from WT of *Arabidopsis thaliana* leaf containing 2.5 µg, and 5 µg of chlorophyll, respectively.

Thy-*stn7*: thylakoid membrane proteins from *stn7* mutant of *Arabidopsis thaliana* leaf containing 5 µg of chlorophyll.

Electrophoresis: 12.5% SDS-Urea-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.