

Anti-SNF1-related protein kinase catalytic subunit alpha KIN10, C-terminal antibody

Catalog: PHY1141A

Product Information

Description:	Rabbit polyclonal antibody	
Background:	In plants, SNF1-related kinase (SnRK1) responds to the availability of	
	carbohydrates as well as to environmental stresses by down-regulating ATP	
	consuming biosynthetic processes, while stimulating energy-generating	
	catabolic reactions through gene expression and post-transcriptional	
	regulation.	
Synonyms:	AKIN10, KIN10, SNF1 KINASE HOMOLOG 10, SNF1-RELATED PROTEIN	
	KINASE 1.1, SNRK1.1	
Immunogen:	KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from	
	Arabidopsis thaliana AKIN10 (AT3G01090).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Immunogen affinity purified	
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\!\! \mathbb{C}$ as supplied.	
	6 months, -20 to -70 $^\circ$ C under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.	
Shipping:	The product is shipped at $4^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected/apparent MW:	61 kDa




Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica rapa*, *Brassica napus*.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only