

Anti-NADH dehydrogenase subunit AT4G16450, mitochondrial antibody

Catalog: PHY0528A

Product Information

Description: Rabbit polyclonal antibody

Background: Complex I is the largest protein complex of the oxidative phosphorylation

system in mitochondrial and it catalyzes NADH-quinone oxidoreduction.

Complex I represents the main entrance site for electrons into the respiratory electron transfer chain. In Arabidopsis, Complex I have at least 49 subunits and

AT4G16450 is one of the subunit.

Synonyms: AT4G16450

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana AT4G16450.

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8 [°]C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4° C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 11 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for


immunization is 100% homologues with the sequence in Oryza sativa,



Spinacia oleracea, Gossypium raimondii, Zea mays, Setaria viridis, Sorghum bicolor, Glycine max, Cucumis sativus, and 80-99% homologues with the sequence in Populus trichocarpa, Solanum tuberosum, Vitis vinifera, Brassica napus, Brassica rapa, Triticum aestivum, Solanum lycopersicum, Nicotiana tabacum, Medicago truncatula, Hordeum vulgare.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.