

## Anti-Small RNA 2'-O-methyltransferase antibody

Catalog: PHY3886S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** HEN1 is an enhancer of hua1 and hua2 tha acts to specify reproductive organ

identities and to repress A gene function. HEN1 also shares AG's non-homeotic

function in controlling floral determinacy.

Synonyms: HEN1, CORYMBOSA 2, CRM2, HUA ENHANCER 1

**Immunogen:** KLH-conjugated synthetic peptide (17 aa from C terminal section) derived from

Arabidopsis thaliana HEN1 (AT4G20910).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability & Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 104 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in *Brassica* 


napus, Brassica rapa, Solanum lycopersicum, Spinacia oleracea,



Cucumis sativus.

For more species homologues information, please contact tech support at <a href="mailto:tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 53 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.