

Anti-Probable UDP-N-acetylglucosamine--peptide N-acetylglucosaminyltransferase SEC, C-terminal antibody

Catalog: PHY1981S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Arabidopsis has two O-GlcNAc transferases (OGTs), SPINDLY (SPY) and SECRET AGENT (SEC). Both enzymes have a domain structure like that of the animal OGTs. OGT activity is required during embryogenesis as loss of both SPY and SEC function results in embryo lethality. In addition to roles in embryogenesis, SPY and SEC have both overlapping and unique roles throughout the life of the plant.
Synonyms:	SEC, SECRET AGENT
Immunogen:	KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> SEC (AT3G04240).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Research Use Only


Note: Optimal dilutions/concentrations should be determined by the end user.

Expected / apparent MW: 110 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica napu*, *Brassica rapa*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.