

Anti-Zinc finger protein ZAT2 antibody

Catalog: PHY7701S

Product Information

Description:	Rabbit polyclonal antibody
Background:	DAZ1 is a target promoter of the male germline-specific transcription factor
	DUO1.
Synonyms:	DAZ1, DUO1-ACTIVATED ZINC FINGER 1, ZAT2
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from
	Arabidopsis thaliana DAZ1 (AT2G17180).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at info@phytoab.com.
Reconstitution:	Reconstitution with 150 μl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ \! \mathbb C$ as supplied.
	6 months, -20 to -70 $^\circ\!\mathrm{C}$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ\!\!\!\!^\circ$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!\mathbb{C}$. Upon receipt, store it immediately at the
	temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	30 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at <u>tech@phytoab.com</u> .

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4° C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only

PhytoAB Inc.