

## Anti-OSCA1.1/1.2, C-terminal antibody

Catalog: PHY7634S

## **Product Information**

Rabbit polyclonal antibody
OSCA1.1/1.2
OSCA1.1/1.2, REDUCED HYPEROSMOLALITY-INDUCED CA2+ INCREASE
1.1 / 1.2
KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from
Arabidopsis thaliana OSCA1.1 (AT4G04340) and CSC1 (AT4G22120).
Lyophilized
150 μg
Serum
Peptide affinity form antibody available upon request at info@phytoab.com.
Reconstitution with 150 µl of sterile water.
"Note: please spin tube briefly prior to opening it to avoid any losses that might
occur from lyophilized material adhering to the cap or sides of the tube".
Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
12 months from date of receipt, -20 to -70 $^\circ\!\mathrm{C}$ as supplied.
6 months, -20 to -70 $^\circ\!{ m C}$ under sterile conditions after reconstitution.
1 month, 2 to 8 $^\circ \!\!\!\!\!^\circ \!\!\!^\circ$ under sterile conditions after reconstitution.
The product is shipped at 4 $^\circ\!\!\mathbb{C}$ . Upon receipt, store it immediately at the
temperature recommended above.

## **Application Information**


<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	88 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Brassica</i>
	napus, Brassica rapa, Populus trichocarpa, and 80-99% homologues
	with the sequence in Goss <i>ypium raimondii, Oryza sativa, Glycine max</i> ,



Vitis vinifera, Spinacia oleracea, Nicotiana tabacum, Zea mays, Panicum virgatum, Hordeum vulgare, Sorghum bicolor, Triticum aestivum, Setaria viridis.

The sequence of the synthetic peptide used for immunization is 87% (13/15) homologues with the sequence in OSCA1.3 (AT1G11960). For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.

**Research Use Only**