

Anti-20S proteasome β subunit 1 antibody

Catalog: PHY2474S

Product Information

Description: Rabbit polyclonal antibody

Background: The plant 20S proteasome β subunit 1 (PBA1) was shown to have

caspase-3-like activity using biotin-DEVD-fmk pull-down and activity assay in

downregulated lines.

Synonyms: PBA1

Immunogen: KLH-conjugated synthetic peptide (19 aa from N terminal section) derived from

Arabidopsis thaliana PBA1 (AT4G31300).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 $^{\circ}$ C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:5000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 25 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Zea mays,


Solanum tuberosum, Panicum virgatum, Sorghum bicolor, Setaria



viridis, Vitis vinifera, Nicotiana tabacum, Solanum lycopersicum, Gossypium raimondii, Brassica rapa, Brassica napus, and 80-99% homologues with the sequence in Triticum aestivum, Oryza sativa, Hordeum vulgare, Spinacia oleracea, Populus trichocarpa, Cucumis sativus, Glycine max, Medicago truncatula.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 55 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.