

## Anti-High-affinity nitrate transporter 2.1, C-terminal antibody

Catalog: PHY7794S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** NRT2.1 is a high-affinity nitrate transporter. It functions as a repressor of lateral

root initiation independently of nitrate uptake.

Synonyms: NRT2.1, ACH1, ATNRT2.1, ATNRT2:1, LATERAL ROOT INITIATION 1, LIN1,

NITRATE TRANSPORTER 2, NITRATE TRANSPORTER 2.1, NITRATE

TRANSPORTER 2:1, NRT2, NRT2:1, NRT2;1AT

**Immunogen:** KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana NRT2.1 (AT1G08090).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 58 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in Brassica



napus, Brassica rapa, Solanum lycopersicum, Solanum tuberosum, Vitis vinifera.

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 41 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.