

## Anti-COP9 signalosome complex subunit 4 antibody

Catalog: PHY7811S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** COP8 is subunit 4 of COP9 signalosome complex. sequence is similar to a

subunit of the 19S regulatory particle of the 26S proteasome.

**Synonyms:** COP8, ATS4, CONSTITUTIVE PHOTOMORPHOGENIC 14, CONSTITUTIVE

PHOTOMORPHOGENIC 8, COP14, COP9 SIGNALOSOME SUBUNIT 4, CSN4, EMB134, EMBRYO DEFECTIVE 134, FUS4, FUSA, FUSCA 4, FUSCA

8

**Immunogen:** KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana COP8 (AT5G42970).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 45 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in Brassica



napus, Brassica rapa and 80-99% homologues with the sequence in Zea mays, Oryza sativa, Panicum virgatum, Setaria viridis, Sorghum bicolor, Triticum aestivum.

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4℃ for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.