

Anti-Calmodulin-binding transcription activator 3 antibody

Catalog: PHY7537A

Product Information

| Description: | Rabbit polyclonal antibody |
|------------------------|--|
| Background: | CAMTA3 is a putative CAM binding transcription factor. It acts in the cold |
| | response pathway. It can bind to and activate the expression of DREB1 genes. |
| Synonyms: | CAMTA3, CALMODULIN-BINDING TRANSCRIPTION ACTIVATOR 3, SIGNAL |
| | RESPONSIVE 1, SR1 |
| Immunogen: | KLH-conjugated synthetic peptide (17 aa from N terminal section) derived from |
| | Arabidopsis thaliana CAMTA3 (AT2G22300). |
| Form: | Lyophilized |
| Quantity: | 150 µg |
| Purification: | Immunogen affinity purified |
| Reconstitution: | Reconstitution with 150 µl of sterile 1XPBS (PH=7.4). |
| | "Note: please spin tube briefly prior to opening it to avoid any losses that might |
| | occur from lyophilized material adhering to the cap or sides of the tube". |
| Stability & | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. |
| Storage: | 12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb{C}$ as supplied. |
| | 6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution. |
| | 1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution. |
| Shipping: | The product is shipped at 4° C. Upon receipt, store it immediately at the |
| | temperature recommended above. |


Application Information

| Recommended Dilution: | Western Blot (1:1000-1:2000) |
|------------------------------|--|
| | Note: Optimal dilutions/concentrations should be determined by the |
| | end user. |
| Expected / apparent MW: | 116 kDa |
| Predicted Reactivity: | For more species homologues information, please contact tech |
| | support at tech@phytoab.com. |

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only