

## Anti-Photosystem I reaction center subunit III antibody

Catalog: PHY4521S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** PSAF is PHOTOSYSTEM 1-F subunit.

**Synonyms:** PSAF, OsPS1-F

**Immunogen:** KLH-conjugated synthetic peptide (15 aa from Central section) derived from

Oryza sativa PSAF (Os03g0778100).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 25 kDa

**Predicted Reactivity:** Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Triticum

aestivum, Hordeum vulgare, and 80-99% homologues with the


sequence in Panicum virgatum, Zea mays, Setaria viridis, Sorghum

bicolor, Spinacia oleracea, Cucumis sativus, Glycine max, Nicotiana



tabacum, Gossypium raimondii, Vitis vinifera, Brassica napus, Brassica rapa, Populus trichocarpa, Solanum lycopersicum, Solanum tuberosum, Arabidopsis thaliana, Medicago truncatula. For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 53 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.