

Anti-Transcription factor PIF7 antibody

Catalog: PHY1952S

Product Information

Description:	Rabbit polyclonal antibody	
Background:	PIF7 is a basic helix-loop-helix (bHLH) phytochrome interacting factor. It	
	interacts specifically with the far-red light-absorbing Pfr form of phyB through a	
	conserved domain called the active phyB binding motif. Upon light exposure,	
	PIF7 rapidly migrates to intranuclear speckles, where it colocalizes with phyB.	
Synonyms:	PIF7, PHYTOCHROME-INTERACTING FACTOR7	
Immunogen:	KLH-conjugated synthetic peptide (17 aa from Central section) derived	
	fromderived from Arabidopsis thaliana PIF7 (AT5G61270).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at info@phytoab.com.	
Reconstitution:	Reconstitution with 150µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.	
	6 months, -20 to -70 $^\circ$ C under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ$ C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information


Recommended Dilution:	Western Blot (1:1000-1:4000);
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	41 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 80-99% homologues with the sequence in <i>Brassica</i>
	napus, Brassica rapa.

lesearch Use Only



For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 53 kDa. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4° C for 1 h. Primary antibody: 1:1000 dilution overnight at 4 °C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). Detection: using chemiluminescence substrate and image were captured

PHY1952S

with CCD camera.

Research Use Only