

Anti-PsaF subunit of Photosystem I antibody

Catalog: PHY0120A

Product Information

Description: Rabbit polyclonal antibody

Background: PsaF participates in efficiency of electron transfer from plastocyanin to P700 (or

cytochrome c553 in algae and cyanobacteria).

Synonyms: PsaF, PHOTOSYSTEM I SUBUNIT F, PSAF

Immunogen: KLH-conjugated synthetic peptide (16 aa from Central section) derived from

Arabidopsis thaliana PsaF (AT1G31330).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 μl of sterile 1XPBS (PH=7.4).

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:5000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected/apparent MW: 24 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in *Brassica* rapa, *Brassica napus*, and 80-99% homologues with the sequence in

Solanum tuberosum, Solanum lycopersicum, Nicotiana tabacum,


Oryza sativa, Zea mays, Setaria viridis, Gossypium raimondii, Triticum



aestivum, Populus trichocarpa.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.