

Anti-Calreticulin-3, C-terminal antibody

Catalog: PHY3621S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Calreticulin (CRT) is an endoplasmic reticulum (ER)-localized chaperone-like
	lectin that plays a crucial role in promoting the folding and maturation of newly
	synthesized glycoproteins and retaining incompletely/mis-folded proteins in the
	ER through its specific binding to monoglucosylated aspariginelinked glycans.
	And the CRTs are including CRT1 (AT1G56340), CRT2 (AT1G09210), CRT3
	(AT1G08450).
Synonyms:	CRT3, CALRETICULIN 3, EBS2, EMS-MUTAGENIZED BRI1 SUPPRESSOR
	2, PRIORITY IN SWEET LIFE 1, PSL1
Immunogen:	KLH-conjugated synthetic peptide (17 aa from C terminal section) derived from
	Arabidopsis thaliana CRT3 (AT1G08450).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
Reconstitution:	Reconstitution with 150µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^{\circ}$ C under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}.$ Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	50 kDa




Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is100% homologues with the sequence in *Brassica napus*, *Brassica rapa*.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 44 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only