

Anti-Protein ABSCISIC ACID-INSENSITIVE 5, C-terminal antibody

Catalog: PHY0773S

Product Information

Description: Rabbit polyclonal antibody

Background: ABI5 is a member of the basic leucine zipper transcription factor family,

involved in ABA signalling during seed maturation and germination. The Arabidopsis abscisic acid (ABA)-insensitive abi5 mutants have pleiotropic

defects in ABA response, including decreased sensitivity to ABA inhibition of

germination and altered expression of some ABA-regulated genes.

Synonyms: ABI5, ABA INSENSITIVE 5, ATABI5, DPBF1, GIA1, GROWTH-INSENSITIVITY

TO ABA 1

Immunogen: KLH-conjugated synthetic peptide (20 aa from C terminal section) derived from

Arabidopsis thaliana ABI5 (AT2G36270).

Form: Lyophilized

Quantity: 150 μg **Purification**: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000), ChIP

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 47 kDa

Research Use Only




Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica rapa*, *Brassica napus*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.