

# Anti-Cucurbit yellow stunting disorder virus (CYSDV) Capsid protein antibody

**Catalog:** NAV1023-001S **Quantity:** 200 μL

# Immunogen Information:

#### Background

Cucurbit yellow stunting disorder virus (CYSDV)

#### Immunogen

KLH-conjugated synthetic peptide (18 aa from Central section) derived from Cucurbit yellow stunting disorder virus CYSDV Capsid protein (Uniprot: Q643W8 NCBI: AAU25942). We also have antibodies for different epitopes from the Capsid protein. Please request at <u>info@nanodiaincs.com</u> or <u>https://www.nanodiaincs.com</u>.

## **Basic Information:**

Purification: SerumPeptide affinity form antibody available upon request at <a href="mailto:info@nanodiaincs.com">info@nanodiaincs.com</a>.Clonality: PolyclonalExpected MW: 29 kDaHost: Rabbit

## **Product Information:**

Form: Lyophilized

#### Reconstitution

Reconstitution with 200  $\mu$ L of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".

#### **Stability & Storage**

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

12 months from date of receipt, -20 to -70  $^\circ\!\mathbb{C}$  as supplied.

6 months, -20 to -70  $^\circ$ C under sterile conditions after reconstitution.

1 month, 2 to  $8^{\circ}$ C under sterile conditions after reconstitution.

#### Shipping

The product is shipped at  $4^{\circ}$ C. Upon receipt, store it immediately at the temperature recommended above.

# **Applications Information:**


Recommended Dilution:WB (1:1000-1:2000)Predicted Reactivity:For more species homologues information, please contact<br/>tech support at info@nanodiaincs.com.

Research Use Only

Nano Diagnostics, LLc.



# **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 46 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.

**Research Use Only** 

Nano Diagnostics, LLc.