

Anti-Endoglucanase 25, C-terminal antibody

Catalog: PHY0742A

Product Information

Description:	Rabbit polyclonal antibody
Background:	KOR1 is a membrane-bound endo-1,4-beta-D-glucanase, involved in cellulose biosynthesis. During cell elongation, KOR1 is associated with Golgi apparatus and early endosome.
Synonyms:	KOR1, ATGH9A1, DEC, DEFECTIVE CYTOKINESIS, GH9A1, GLYCOSYL HYDROLASE 9A1, IRREGULAR XYLEM 2, IRX2, KOR, KORRIGAN, KORRIGAN 1, RADIALLY SWOLLEN 2, RSW2, TSD1, TUMOROUS SHOOT DEVELOPMENT 1
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> KOR1 (AT5G49720).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen Affinity Purified
Reconstitution:	Reconstitution with 150 µl of sterile 1×PBS (PH=7.4). "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	69 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in

Research Use Only

Brassica rapa, Brassica napus, Zea mays, Cucumis sativus, Gossypium raimondii, Spinacia oleracea, Solanum tuberosum, Solanum lycopersicum.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 54 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.