

Anti-PIP1;1, PIP1;2 aquaporins antibody

Catalog: PHY0051S

Product Information

Description: Rabbit polyclonal antibody

Background: PIPs proteins are aquaporins which selectively conduct water molecules in and

out of the cell, while preventing the passage of ions and other solutes. PIPs are

also known as water channels, aquaporins are integral membrane pore proteins. Some of them, known as aquaglyceroporins, also transport other

small uncharged solutes, such as glycerol, CO2, ammonia and urea across the

membrane, depending on the size of the pore.

Synonyms: PIP1;1/2

Immunogen: KLH-conjugated synthetic peptide of PIP1;1, PIP1;2 derived from *Arabidopsis*

thaliana AT3G61430, AT2G45960.

Form: Lyophilized

Quantity: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected/apparent MW: 31 / 28 kDa



Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among 25 analyzed species, the sequence of the synthetic peptide

used for immunization is 100% homologues with the sequence in

Oryza sativa Japonica Group, Zea mays, Sorghum bicolor, Nicotiana

tabacum, Vitis vinifera, Brassica napus.

The sequence of the synthetic peptide used for immunization is 93%


homologues with the sequence in PIP1;3 (AT1G01620), PIP1;4

(AT4G00430), PIP1;5 (AT4G23400).

For more species homologues information, please contact tech

support at tech@phytoab.com.

Application Example



Lane 1: 20 µg total protein from Arabidopsis thaliana leaf.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

PHY0051S Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.