

Anti-Os07g0673900, C-terminal antibody

Catalog: PHY4555S

Product Information

Rabbit polyclonal antibody
OsHIGD2 is Hypoxia-Induced Gene Domain 2.
OsHIGD2, HIGD2
KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from
<i>Oryza sativa</i> OsHIGD2 (Os07g0673900).
Lyophilized
150 µg
Serum
Peptide affinity form antibody available upon request at info@phytoab.com.
Reconstitution with 150 μ l of sterile water.
"Note: please spin tube briefly prior to opening it to avoid any losses that might
occur from lyophilized material adhering to the cap or sides of the tube".
Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
12 months from date of receipt, -20 to -70 $^\circ \! \mathbb{C}$ as supplied.
6 months, -20 to -70 $^\circ\!{ m C}$ under sterile conditions after reconstitution.
1 month, 2 to 8 $^\circ\!\!\!\!^\circ$ under sterile conditions after reconstitution.
The product is shipped at 4 $^\circ\!{ m C}$. Upon receipt, store it immediately at the
temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	11 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at <u>tech@phytoab.com</u> .

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only

PhytoAB Inc.