

Anti-Os01g0813100 protein, C-terminal antibody

Catalog: PHY4563S

Product Information

Description: Rabbit polyclonal antibody

Background: Os01g0813100 **Synonyms:** Os01g0813100

Immunogen: KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Oryza sativa Os01g0813100.

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 37 kDa

Predicted Reactivity: Among species analyzed, the seugence of the synthetic peptide used

for immunization is 80-99% homologues eith the sequence in


Panicum virgatum, Solanum lycopersicum, Nicotiana tabacum, Vitis vinifera, Gossypium raimondii, Cucumis sativus, Hordeum vulgare, Triticum aestivum, Solanum tuberosum, Zea mays, Sorghum bicolor,



Populus trichocarpa.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.