

## Anti-D1 subunit of Photosystem II, C-terminal antibody

Catalog: PHY3191S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** The D1 protein of Photosystem II (PSII), encoded by the *PsbA* genes, is an

indispensable component of oxygenic photosynthesis. Due to strongly oxidative

chemistry of PSII water splitting, the D1 protein is prone to constant

photodamage requiring its replacement, whereas most of the other PSII

subunits remain ordinarily undamaged.

Synonyms: PsbA, 32 kDa thylakoid membrane protein, Photosystem II protein D1

Immunogen: KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from

Arabidopsis thaliana PsbA (ATCG00020).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at  $4^{\circ}$ C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 39 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in Oryza



sativa, Medicago truncatula, Synechococcus elongatus PCC 7942, Vitis vinifera, Gossypium raimondii, Triticum aestivum, Synechocystis sp. PCC 6803, Glycine max, Brassica napus, Populus trichocarpa, Hordeum vulgare, Chlamydomonas reinhardtii, Zea mays, Nicotiana tabacum, Solanum tuberosum, Brassica rapa, Cucumis sativus, Panicum virgatum, Spinacia oleracea, Setaria viridis.

For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng and 15 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4℃ for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with

CCD camera.