

# Anti-RbcL subunit of RuBisCO antibody

Catalog: PHY0104A

### **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** Ribulose-1,5-bisphosphate carboxylase/oxygenase commonly known by the

abbreviation RuBisCO, is an enzyme involved in the first major step of carbon fixation, a process by which atmospheric carbon dioxide is converted by plants to energy-rich molecules such as glucose. In chemical terms, it catalyzes the

carboxylation of ribulose-1,5-bisphosphate (also known as RuBP). It is

probably the most abundant enzyme on Earth.

The enzyme usually consists of two types of protein subunit, called the large

chain (RbcL) and the small chain (RbcS).

**Synonyms:** RbcL, Ribulose-1,5-bisphosphate carboxylase, oxygenase

**Immunogen:** KLH-conjugated synthetic peptide (16 aa from Central section) derived from

Arabidopsis thaliana RbcL (ATCG00490).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen Affinity Purified

**Reconstitution:** Reconstitution with 150  $\mu$ l of sterile 1 $\times$  PBS (PH=7.4).

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:5000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 53 kDa

Research Use Only




### **Predicted Reactivity:**

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in Solanum tuberosum, Zea mays, Sorghum bicolor, Gossypium raimondii, Vitis vinifera, Brassica rapa, Leymus chinensis, Setaria viridis, Oryza sativa, Nicotiana tabacum, Medicago truncatula, Populus trichocarpa, Glycine max, Cucumis sativus, Hordeum vulgare, Brassica napus, Solanum lycopersicum, Panicum virgatum, Triticum aestivum, Spinacia oleracea.

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

### Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.