

Anti-Protein SUPPRESSOR OF npr1-1, CONSTITUTIVE 1 antibody

Catalog: PHY1863A

Product Information

Description:	Rabbit polyclonal antibody
Background:	SNC1 is a Toll Interleukin1 receptor-nucleotide binding-Leu-rich repeat-type resistance gene (TIR-NB-LRR-type) involved in the salicylic acid-dependent defense response pathway.
Synonyms:	SNC1, BAL, BALL, SUPPRESSOR OF NPR1-1, CONSTITUTIVE 1
Immunogen:	KLH-conjugated synthetic peptide (18 aa from Central section) derived from <i>Arabidopsis thaliana</i> SNC1 (AT4G16890).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	162 kDa
Confirmed Reactivity:	Coming soon
Predicted Reactivity:	The sequence of the synthetic peptide used for immunization is 83% homologues with the sequence in BAR1 (AT5G18360).

Research Use Only

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 5 ng and 15 ng recombinant protein containing the peptide for immunization and having a molecular mass of 55 kDa.

Electrophoresis: 12% SDS-PAGE.

Transfer: blotting to NC (nitrocellulose) membrane for 1h.

Blocking: 5% skim milk at RT or 4°C for 1h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.