

# Anti-Ribulose bisphosphate carboxylase large chain, N-terminal antibody

Catalog: PHY5236A

#### **Product Information**

**Description:** Rabbit polyclonal antibody

Background: cbbL
Synonyms: cbbL

**Immunogen:** KLH-conjugated synthetic peptide (11 aa from N terminal section)

Synechocystis sp. PCC 6803 cbbL (slr0009).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 52 / 53 kDa

Confirmed Reactivity: Synechocystis sp. PCC 6803

**Predicted Reactivity:** Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in

Synechococcus elongatus PCC 7942.

For more species homologues information, please contact tech



#### support at tech@phytoab.com.

## **Application Example**

PCC 6803-TP: 31 µg and 62 µg total protein from Synechocystis sp. PCC 6803,

respectively.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at  $4^{\circ}$ C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.