

# Anti-Histone acetyltransferase GCN5 antibody

Catalog: PHY0852A

## Product Information

<b>Description:</b>	Rabbit polyclonal antibody
<b>Background:</b>	Histone acetyltransferase GCN5 plays a role in the determination of the embryonic root-shoot axis. It is also required to regulate the floral meristem activity by modulating the extent of expression of WUS and AG. HAG1 is specific for histone H3 lysine 14.
<b>Synonyms:</b>	GCN5, BGT, BIG TOP, GENERAL CONTROL NONDEREPRESSIBLE 5, HAC3, HAG01, HAG1, HAT1, HISTONE ACETYLTRANSFERASE 1, HISTONE ACETYLTRANSFERASE OF THE GNAT FAMILY 1
<b>Immunogen:</b>	KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> GCN5 (AT3G54610).
<b>Form:</b>	Lyophilized
<b>Quantity:</b>	150 µg
<b>Purification:</b>	Immunogen affinity purified
<b>Reconstitution:</b>	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
<b>Stability &amp; Storage:</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
<b>Shipping:</b>	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

## Application Information

<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
<b>Expected / apparent MW:</b>	63 kDa
<b>Confirmed Reactivity:</b>	Coming soon


Research Use Only

**Predicted Reactivity:**

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Gossypium raimondii*, *Brassica napus*, *Brassica rapa*, *Medicago truncatula*.

For more species homologues information, please contact tech support at [tech@phytoab.com](mailto:tech@phytoab.com).

**Application Example**



Recom: 5 ng and 15 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

**Electrophoresis:** 12% SDS-PAGE.

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.