

# Anti-c-Myc antibody

Catalog: PHY5014

### **Product Information**

**Description:** Mouse monoclonal antibody

**Synonyms**: c-Myc

**Immunogen:** Synthetic peptide of fragment (408-420 aa) c-Myc.

Form: Lyophilized

**Quantity:** 150 μg

**Purification:** Protein A purified

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the


end user.

**Confirmed Reactivity:** Anti-c-Mycrecognizesc-Myc-tagged fusion proteins.



### **Application Example**

### Example 1



Recom: 2.5 ng and 5 ng c-Myc-tagged fusion protein, respectively and having

a molecular mass of 56 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4 ☐ for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.


Secondary antibody: 1:5000 dilution using Goat Anti-Mouse IgGH&L

(HRP) (Cat# PHY6006).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.

### Example 2



TP: 30 µg total protein from WT of Arabidopsis thaliana.

Myc-TP: 30 µg total protein from transgenic Arabidopsis thaliana, containing c-

Myc-tagged fusion protein.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:5000 dilution using Goat Anti-Mouse IgGH&L

(HRP) (Cat# PHY6006).

**Detection:** using chemiluminescence substrate and image were captured with

CCD camera.