

# Anti-PLASTID TRANSCRIPTIONALLY ACTIVE 3 antibody

Catalog: PHY0391A

## Product Information

<b>Description:</b>	Rabbit polyclonal antibody
<b>Background:</b>	In chloroplasts, transcription of plastid genes is mediated by two types of RNA polymerase: plastid-encoded RNA polymerase (PEP) and nuclearencoded RNA polymerase (NEP). Transcription in plastids is also mediated by a number of nuclear-encoded factors in addition to PEP and NEP. In the insoluble RNA polymerase preparation samples, a total of 18 components named as pTACs (pTAC1 to pTAC18) were identified. pTAC3 (AT3G04260) is one of the components associated with PEP complex.
<b>Synonyms:</b>	pTAC3, PDE324, PIGMENT DEFECTIVE 324, PLASTID TRANSCRIPTIONALLY ACTIVE 3
<b>Immunogen:</b>	KLH-conjugated synthetic peptide (15 aa from central section) of derived from <i>Arabidopsis thaliana</i> pTAC3 (AT3G04260).
<b>Form:</b>	Lyophilized
<b>Quantity:</b>	150 µg
<b>Purification:</b>	Immunogen affinity purified
<b>Reconstitution:</b>	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
<b>Stability &amp; Storage:</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
<b>Shipping:</b>	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

## Application Information

<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
------------------------------	--

Research Use Only


**Expected / apparent MW:** 103 kDa

**Confirmed Reactivity:** *Arabidopsis thaliana*

**Predicted Reactivity:** Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Glycine max*, *Brassica rapa*, *Panicum virgatum*, *Brassica napus*, *Oryza sativa*, *Gossypium raimondii*, *Spinacia oleracea*, *Setaria viridis*, *Solanum lycopersicum*, *Solanum tuberosum*, *Populus trichocarpa*, *Zea mays*, *Sorghum bicolor*, *Hordeum vulgare subsp. vulgare*, *Vitis vinifera*, *Triticum aestivum*, *Nicotiana tabacum*, and 80-99% homologues with the sequence in *Cucumis sativus*, *Medicago truncatula*.

For more species homologues information, please contact tech support at [tech@phytoab.com](mailto:tech@phytoab.com).

## Application Example



Chl: 10  $\mu$ l total chloroplast protein from *Arabidopsis thaliana*.

**Electrophoresis:** 10% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:2000 dilution overnight at 4°C.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.