

Anti-PsaO subunit of photosystem I, C-terminal antibody

Catalog: PHY0648A

Product Information

Description:	Rabbit polyclonal antibody
Background:	PsaO has two transmembrane helices and a calculated mass of 10104 Da. PsaO is involved in the balancing of excitation energy between the two photosystems I (PSI) and II (PSII).
Synonyms:	PsaO, Photosystem I subunit O, PSI-O, PHOTOSYSTEM I SUBUNIT O, PSAO
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> PsaO (AT1G08380).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability &Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	15 / 11 kDa
Confirmed Reactivity:	<i>Arabidopsis thaliana</i>
Predicted Reactivity:	Among species analyzed , the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in

Research Use Only

Populus trichocarpa, Brassica napus, Vitis vinifera, Panicum virgatum, Cucumis sativus, Solanum lycopersicum, Brassica rapa, Solanum tuberosum, Nicotiana tabacum, Triticum aestivum, Hordeum vulgare, Zea mays, Setaria viridis, Sorghum bicolor, Glycine max, Medicago truncatula, and 80-99% homologues with the sequence in *Spinacia oleracea, Gossypium raimondii, Oryza sativa*.
For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Thy-WT: thylakoid membrane protein from WT of *Arabidopsis thaliana* containing 0.25 μ g of chlorophyll.

Thy-PS I : thylakoid membrane protein from an *Arabidopsis* mutant with low accumulation of PS I containing 0.25 μ g of chlorophyll.

Electrophoresis: Tricine-SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.