

Anti-Ferredoxin--NADP reductase, leaf isozyme 1, chloroplastic, N-terminal antibody

Catalog: PHY2232S

Product Information

Description: Rabbit polyclonal antibody

Background: Two distinct ferredoxin-NADP(+)-oxidoreductase (FNR) isoforms were found in

chloroplasts of Arabidopsis thaliana, FNR-1 (AT5G66190) and FNR-2

(AT1G20020). The FNR proteins are present in both chloroplast stroma and thylakoid membranes in chloroplasts but are more abundant in the stroma.

Synonyms: FNR1, ATLFNR1, FERREDOXIN-NADP(+)-OXIDOREDUCTASE 1, LEAF FNR

1, LEAF-TYPE CHLOROPLAST-TARGETED FNR 1, LFNR1

Immunogen: KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from

Arabidopsis thaliana FNR1 (AT5G66190).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 40 / 34 kDa




Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: For more species homologues information, please contact tech

support at tech@phytoab.com.

Application Example



Str: 12 µg stromal protein from *Arabidopsis thaliana*.

Thy: thylakoid membrane protein from Arabidopsis thaliana containing

5 µg of chlorophyll.

Chl: 10 µl total chloroplast protein from Arabidopsis thaliana.

TP: 18 µg total protein from *Arabidopsis thaliana*.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L (HRP) (Cat# PHY6000)

Detection: using chemiluminescence substrate and image were

captured with CCD camera.