

# Anti-ATP-dependent Clp protease proteolytic subunit 3, chloroplastic antibody

Catalog: PHY2069A

## Product Information

<b>Description:</b>	Rabbit polyclonal antibody
<b>Background:</b>	Caseinolytic protease (Clp) is a major protease system in plant cells and functions to remove mis-folded, damaged, short-lived regulatory, and otherwise unneeded proteins in plastids to maintain protein homeostasis. CLPP3 is one of several nuclear-encoded ClpPs (caseinolytic protease).
<b>Synonyms:</b>	CLPP3, CLP PROTEASE PROTEOLYTIC SUBUNIT 3, NCLPP3
<b>Immunogen:</b>	KLH-conjugated synthetic peptide of CLPP3 (16 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> (AT1G66670).
<b>Form:</b>	Lyophilized
<b>Quantity:</b>	150 µg
<b>Purification:</b>	Immunogen affinity purified
<b>Reconstitution:</b>	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
<b>Stability &amp; Storage:</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
<b>Shipping:</b>	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

## Application Information


<b>Recommended Dilution:</b>	Western Blot(1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
<b>Expected / apparent MW:</b>	34 / 26 kDa
<b>Confirmed Reactivity:</b>	<i>Arabidopsis thaliana</i>
<b>Predicted Reactivity:</b>	Among species analyzed, the sequence of the synthetic peptide used

Research Use Only

for immunization is 80-99% homologues with the sequence in *Brassica napus*, *Brassica rapa*, *Gossypium raimondii*.

For more species homologues information, please contact tech support at [tech@phytoab.com](mailto:tech@phytoab.com).

## Application Example



Str: 8.75 µg and 17.5 µg stromal protein from *Arabidopsis thaliana*, respectively.

**Electrophoresis:** 15% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:2000 dilution overnight at 4°C.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.