

Anti-Homogentisate phytyltransferase 1, chloroplastic antibody

Catalog: PHY2797A

Product Information

Description:	Rabbit polyclonal antibody	
Background:	VTE2 is involved in tocopherol biosynthesis. It has impact on seed longevity	
	and plays a role in the adaptation to low temperature stress, notably phloem	
	loading.	
Synonyms:	VTE2, ATHPT, HOMOGENTISATE PHYTYLTRANSFERASE,	
	HOMOGENTISATE PHYTYLTRANSFERASE 1, HPT1, TPT1, VITAMIN E 2	
Immunogen:	KLH-conjugated synthetic peptide (18 aa from N terminal section) derived from	
	Arabidopsis thaliana VTE2 (AT2G18950).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Immunogen affinity purified	
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!$	
	6 months, -20 to -70 $^\circ C$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.	
Shipping:	The product is shipped at $4^\circ C$. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	44 kDa
Confirmed Reactivity:	Arabidopsis thaliana
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide
	used for immunization is 100% homologues with the sequence in


PhytoAB Inc.



Solanum tuberosum, and 80-99% homologues with the sequence in Brassica napus, Brassica rapa. For more species homologues information, please contact tech

support at tech@phytoab.com.

Application Example



Chl: 8 µg total chloroplast protein from Arabidopsis thaliana. TP: 30 µg total protein from Arabidopsis thaliana. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h. Blocking: 5% skim milk at RT or 4°C for 1 h. Primary antibody: 1:1000 dilution overnight at 4°C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). Detection: using chemiluminescence substrate and image were captured with CCD camera.

PHY2797A

Research Use Onl