

Anti-CRS2-associated factor 2, chloroplastic, C-terminal antibody

Catalog: PHY3964S

Product Information

| Description: | Rabbit polyclonal antibody | |
|-----------------|--|--|
| Background: | CAF2 promotes the splicing of chloroplast group II introns. | |
| Synonyms: | CAF2, ARABIDOPSIS THALIANA HOMOLOG OF MAIZE CAF2, ATCAF2, | |
| | CHLOROPLAST RNA SPLICING2-ASSOCIATED FACTOR 2 | |
| Immunogen: | KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from | |
| | Arabidopsis thaliana CAF2 (AT1G23400). | |
| Form: | Lyophilized | |
| Quantity: | 150 μg | |
| Purification: | Serum | |
| | Peptide affinity form antibody available upon request at info@phytoab.com. | |
| Reconstitution: | Reconstitution with 150 µl of sterile water. | |
| | "Note: please spin tube briefly prior to opening it to avoid any losses that might | |
| | occur from lyophilized material adhering to the cap or sides of the tube". | |
| Stability & | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. | |
| Storage: | 12 months from date of receipt, -20 to -70 $^\circ \! \mathbb C$ as supplied. | |
| | 6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution. | |
| | 1 month, 2 to 8 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution. | |
| Shipping: | The product is shipped at 4 $^\circ\!\!{ m C}$. Upon receipt, store it immediately at the | |
| | temperature recommended above. | |
| | | |


Application Information

| Recommended Dilution: | Western Blot (1:1000-1:2000) |
|-------------------------|--|
| | Note: Optimal dilutions/concentrations should be determined by the |
| | end user. |
| Expected / apparent MW: | 64 kDa |
| Predicted Reactivity: | For more species homologues information, please contact tech |
| | support at tech@phytoab.com. |

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 52 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only

PhytoAB Inc.