

Anti-Ethylene insensitive 2 antibody

Catalog: PHY3837S

Product Information

Description: Rabbit polyclonal antibody

Background: EIN2 is a positive regulator in ethylene signaling in Arabidopsis. It acts

downstream of CTR1. It positively regulates ORE1 and negatively regulates

mir164A, B, C to regulate leaf senescence.

Synonyms: EIN2, ATEIN2, CKR1, CYTOKININ RESISTANT 1, ENHANCED RESPONSE

TO ABA3, ERA3, ETHYLENE INSENSITIVE 2, ORE2, ORE3, ORESARA 2,

ORESARA 3, PIR2

Immunogen: KLH-conjugated synthetic peptide (16 aa from Central section) derived from

Arabidopsis thaliana EIN2 (AT5G03280).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected/apparent MW: 141 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in Brassica

rapa, Brassica napus.



For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 44 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit lgG H&L (HRP)

(Cat# PHY6000)

Detection: using chemiluminescence substrate and image were captured with

CCD camera.