

Anti-Probable galacturonosyltransferase 7 antibody

Catalog: PHY1463S

Product Information

Description:	Rabbit polyclonal antibody	
Background:	GAUT7 and Galacturonosyltransferase (GAUT)1 (AT3G61130) are the core of	
	a plant cell wall pectin biosynthetic homogalacturonan:Galacturonosyltran-	
	sferase complex. It may be involved in pectin biosynthesis.	
Synonyms:	GAUT7, GALACTURONOSYLTRANSFERASE 7, JS33, LGT7, LIKE	
	GLYCOSYL TRANSFERASE 7	
Immunogen:	KLH-conjugated synthetic peptide (22 aa from N terminal section) derived from	
	Arabidopsis thaliana GAUT7 (AT2G38650).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .	
Reconstitution:	Reconstitution with 150 μl of ste <mark>rile water.</mark>	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb C$ as supplied.	
	6 months, -20 to -70 $^\circ$ C under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ\!\mathrm{C}$ under sterile conditions after reconstitution.	
Shipping:	The product is shipped at $4^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	70 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 80-99% homologues with the sequence in
	Brassica napus, Brassica rapa.

Research Use Only



For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only