

Anti-ATP-dependent Clp protease proteolytic subunit 6, chloroplastic, C-terminal antibody

Catalog: PHY2063A

Product Information

Description: Rabbit polyclonal antibody

Background: CLPP6 is one of several nuclear-encoded ClpPs (caseinolytic protease). It

cleaves peptides in various proteins in a process that requires ATP hydrolysis, and has a chymotrypsin-like activity. It plays a major role in the degradation of

misfolded proteins.

Synonyms: CLPP6, CLP PROTEASE PROTEOLYTIC SUBUNIT 6, NCLPP1, NCLPP6,

NUCLEAR-ENCODED CLPP 1

Immunogen: KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana CLPP6 (AT1G11750).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 31 kDa


Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 80-99% homologues with the sequence in



Gossypium raimondii, Spinacia oleracea, Glycine max, Populus trichocarpa, Vitis vinifera, Brassica rapa, Nicotiana tabacum, Medicago truncatula, Brassica napus, Medicago truncatula. For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 44 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° °C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.