

## Anti-Transcription termination factor MTERF9, chloroplastic, C-terminal antibody

Catalog: PHY2263S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** MTERF9 is a mitochondrial transcription termination factor (mTERF) family

protein. Loss of MTERF9 results in defective chloroplast development, which is likely to cause paleness, stunted growth and reduced mesophyll cell numbers.

Synonyms: MTERF9

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana MTERF9 (AT5G55580).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 57 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in Brassica



napus, Brassica rapa, and 80-99% homologues with the sequence in Populus trichocarpa, Zea mays, Panicum virgatum, Gossypium raimondii, Cucumis sativus, Setaria viridis, Sorghum bicolor, Glycine max, Oryza sativa, Triticum aestivum, Hordeum vulgare, Vitis vinifera.

For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

for immunization and having a molecular mass of 46 kDa.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.