

## Anti-ATP synthase subunit 4, mitochondrial antibody

Catalog: PHY1129A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** Mitochondrial F0F1-ATP synthase is also called Complex V and it synthesis

ATP from ADP and Pi using the proton motive force created by respiratory

electron transport. ATP4 (ATMG00640) is a subunit of mitochondrial F0F1-ATP

synthase in Arabidopsis.

**Synonyms:** ATP4, ORF25

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana ATP4 (ATMG00640).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 22 / 20 kDa

Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Brassica



rapa, Brassica napus, and 80-99% homologues with the sequence in Glycine max, Vitis vinifera, Cucumis sativus, Medicago truncatula, Gossypium raimondii, Solanum tuberosum, Zea mays, Panicum virgatum, Nicotiana tabacum, Solanum lycopersicum, Oryza sativa, Triticum aestivum, Hordeum vulgare, Spinacia oleracea.

For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**

Mito: 9 µg mitochondria protein from Arabidopsis thaliana.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:2000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat#

PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD

camera.