

Anti-Os06g0473200 antibody

Catalog: PHY4153S

Product Information

| Description: | Rabbit polyclonal antibody | |
|------------------------|--|--|
| Background: | Similar to NPK2. (Os06t0473200-01); Similar to Mitogen-activated protein | |
| | kinase kinase 3. (Os06t0473200-02) | |
| Synonyms: | Os06g0473200, OsMKK3 | |
| Immunogen: | KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from | |
| | <i>Oryza sativa</i> Os06g0473200. | |
| Form: | Lyophilized | |
| Quantity: | 50 µg | |
| Purification: | Serum | |
| Reconstitution: | Reconstitution with 50µl of sterile water. | |
| | "Note: please spin tube briefly prior to opening it to avoid any losses that might | |
| | occur from lyophilized material adhering to the cap or sides of the tube". | |
| Stability & | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. | |
| Storage: | 12 months from date of receipt, -20 to -70 $^\circ\!\mathrm{C}$ as supplied. | |
| | 6 months, -20 to -70 $^\circ \mathrm{C}$ under sterile conditions after reconstitution. | |
| | 1 month, 2 to 8 $^\circ\!\!\mathbb{C}$ under sterile conditions after reconstitution. | |
| Shipping: | The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the | |
| | temperature recommended above. | |

Application Information

| Recommended Dilution: | Western Blot (1:1000-1:2000) |
|------------------------------|---|
| | Note: Optimal dilutions/concentrations should be determined by the |
| | end user. |
| Expected / apparent MW: | 58 kDa |
| Confirmed Reactivity: | Oryza sativa |
| Predicted Reactivity: | Among species analyzed, the sequence of the synthetic peptide used |
| | for immunization is 80-99% homologues with the sequence in |
| | Triticum aes <mark>tivum,</mark> Hordeum vulgare, Panicum virgatum, Setaria |
| | viridis, Sorghum bicolor, Zea mays. |

Research Use Only



For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example

OS-TP: total protein from *Oryza sativa*.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only