

Anti-WRKY68 antibody

Catalog: PHY4270S

Product Information

Description:	Rabbit polyclonal antibody	
Background:	WRKY transcription factor 68. (Os04t0605100-01)	
Synonyms:	WRKY68 / OsWRKY68	
Immunogen:	KLH-conjugated synthetic peptide (13 aa from Central section) derived from	
	<i>Oryza sativa</i> WRKY68 (Os04g0605100).	
Form:	Lyophilized	
Quantity:	50 µg	
Purification:	Serum	
Reconstitution:	Reconstitution with 50µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &		
Stability & Storage:	occur from lyophilized material adhering to the cap or sides of the tube".	
	occur from lyophilized material adhering to the cap or sides of the tube". Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
	occur from lyophilized material adhering to the cap or sides of the tube". Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied.	
	occur from lyophilized material adhering to the cap or sides of the tube". Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution.	

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	32 kDa
Confirmed Reactivity:	Oryza sativa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Zea</i>
	mays, Sorghum bicolor, and 80-99% homologues with the sequence
	in Triticum aestivum, Hordeum vulgare, Setaria viridis, Panicum
	virgatum.



For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example

OS-TP: total protein from *Oryza sativa*.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only