

Anti-Chloroplast outer envelope membrane translocon complex OEP75 protein antibody

Catalog: PHY2973A

Product Information

Description:	Rabbit polyclonal antibody
Background:	Toc75 is the component of the translocon outer membrane (TOC) complex. It forms the outer envelope translocation channel (beta-barrel). It plays a role in preprotein conductance.
Synonyms:	TOC75-III, 1-MAR, MAR1, MODIFIER OF ARG1 1, TOC75, TRANSLOCON AT THE OUTER ENVELOPE MEMBRANE OF CHLOROPLASTS 75-III
Immunogen:	KLH-conjugated synthetic peptide of TOC75-III (15 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> (AT3G46740).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	89 / 70 kDa
Confirmed Reactivity:	<i>Arabidopsis thaliana</i>
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used


Research Use Only

for immunization is 80-99% homologues with the sequence in *Brassica rapa*, *Brassica napus*, *Populus trichocarpa*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example

Chl: 5 μ l and 10 μ l total chloroplast protein from *Arabidopsis thaliana*, respectively.



Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.