

Anti-30S ribosomal protein S8, chloroplastic, C-terminal antibody

Catalog: PHY0429A

Product Information

Description: Rabbit polyclonal antibody

Background: RPS8 is one of the primary rRNA binding proteins. It binds directly to 16S rRNA

central domain where it helps coordinate assembly of the platform of the 30S

subunit.

Synonyms: RPS8, RIBOSOMAL PROTEIN S8

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana RPS8 (ATCG00770).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 15 kDa

Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Brassica



rapa, Brassica napus, and 80-99% homologues with the sequence in Triticum aestivum, Oryza sativa Japonica Group, Setaria viridis, Zea mays, Panicum virgatum, Hordeum vulgare subsp. spontaneum, Sorghum bicolor, Glycine max, Medicago truncatula, Populus trichocarpa, Vitis vinifera, Cucumis sativus, Spinacia oleracea, Nicotiana tabacum, Solanum lycopersicum.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example

Chl: 80 µg total chloroplast protein from *Arabidopsis thaliana*.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.

kDa M

10