

Anti-Cytochrome b6-f complex subunit 5, C-terminal antibody

Catalog: PHY1229S

Product Information

Description:	Rabbit polyclonal antibody
Background:	PetG is cytochrome b6-f complex subunit V which is the component of the cytochrome b6-f complex, which mediates electron transfer between photosystem II (PSII) and photosystem I (PSI), cyclic electron flow around PSI, and state transitions. It is required for either the stability or assembly of the cytochrome b6-f complex.
Synonyms:	PetG
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> PetG (ATCG00600).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	4 kDa

Research Use Only


Confirmed Reactivity:

Arabidopsis thaliana

Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica napus*, and 80-99% homologues with the sequence in *Cucumis sativus*, *Triticum aestivum*, *Physcomitrium patens*, *Gossypium raimondii*, *Oryza sativa*, *Medicago truncatula*, *Spinacia oleracea*, *Brassica rapa*, *Panicum virgatum*, *Glycine max*, *Solanum tuberosum*. For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



captured with CCD camera.

Thy-WT: thylakoid membrane protein from WT of *Arabidopsis thaliana* leaf containing 2 µg and 4 µg of chlorophyll, respectively.

Thy-cyt b6f: thylakoid membrane protein from *Arabidopsis* mutant with low accumulation of Cyt b6f complex containing 4 µg of chlorophyll.

Electrophoresis: Tricine-SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were