

Anti-Peroxisomal membrane protein PEX14, N-terminal antibody

Catalog: PHY0168S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Pex14p has a protein transporter activity and is involved in protein targeting into peroxisome.
Synonyms:	Pex14, Pex14p, ATPED2, ATPEX14, PED2, PEROXIN 14, PEROXISOME DEFECTIVE 2, PEX14
Immunogen:	KLH-conjugated synthetic peptide (18 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> Pex14 (AT5G62810).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	56 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Solanum tuberosum</i> , <i>Brassica napus</i> , <i>Brassica rapa</i> , <i>Nicotiana tabacum</i> ,

Research Use Only

Gossypium raimondii, *Solanum lycopersicum*, *Cucumis sativus*,
Physcomitrium patens, and 80-99% homologues with the sequence in
Glycine max, *Vitis vinifera*, *Populus trichocarpa*, *Medicago truncatula*,
Triticum aestivum, *Hordeum vulgare*, *Oryza sativa*, *Spinacia oleracea*,
Setaria viridis.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.