

Anti-Protein argonaute 4 antibody

Catalog: PHY0202S

Product Information

Description:	Rabbit polyclonal antibody
Background:	AGO4 is a member of a class of PAZ/PIWI domain containing proteins
	involved in siRNA mediated gene silencing. Loss of function mutations have
	reduced site specific CpNpG and CpHpH methylation, abnormal
	ovule/megagametophyte develoment and increased susceptibility to bacterial
	pathogens including Tobacco rattle virus.
Synonyms:	AGO4, ARGONAUTE 4, OCP11, OVEREXPRESSOR OF CATIONIC
	PEROXIDASE 11
Immunogen:	KLH-conjugated synthetic peptide (17 aa from N terminal section) derived from
	Arabidopsis thaliana AGO4 (AT2G27040).
Form:	Lyophilized
Quantity:	150 μg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
Reconstitution:	Reconstitution with 150µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^{\circ}$ C under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!{ m C}$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	103 kDa




Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica napus, Brassica rapa.* For more species homologues information, please contact tech

support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 47 kDa. **Electrophoresis:** 12% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4°C. **Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only