

## Anti-Chaperonin-like RbcX protein 2, chloroplastic antibody

Catalog: PHY7252S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** RBCX2 is a chloroplast stromal localized RbcX protein that acts as a

chaperone in the folding of Rubisco.

**Synonyms:** RBCX2, ATRBCX2, HOMOLOGUE OF CYANOBACTERIAL RBCX 2,

**Immunogen:** KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana RBCX2 (AT5G19855).

Form: Lyophilized

Quantity: $150 \mu g$ Purification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70 □ as supplied.

6 months, -20 to -70 under sterile conditions after reconstitution.

1 month, 2 to 8□ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4□. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 23 kDa

**Predicted Reactivity:** Among species analyzed, the sequence of the synthetic peptide

used for immunization is 80-99% homologues with the sequence in


Brassica napus, Brassica rapa, Zea mays, Triticum aestivum, Hordeum vulgare, Oryza sativa, Sorghum bicolor, Populus



trichocarpa, Gossypium raimondii.

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 40 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.