

## Anti-Tocopherol cyclase, chloroplastic antibody

Catalog: PHY3414A

## **Product Information**

Description:	Rabbit polyclonal antibody	
Background:	Tocopherol cyclase involved in tocopherol (vitamin E) synthesis. It plays a role	
	in the adaptation to low temperature stress, notably phloem loading.	
Synonyms:	VTE1, ATSDX1, SUCROSE EXPORT DEFECTIVE 1, VITAMIN E DEFICIENT	
	1	
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from	
	Arabidopsis thaliana VTE1 (AT4G32770).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .	
<b>Reconstitution:</b>	Reconstitution with 150 µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ\mathrm{C}$ as supplied.	
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!{ m C}$ . Upon receipt, store it immediately at the	
	temperature recommended above.	
Application Information		

<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	55 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide
	used for immunization is 100% homologues with the sequence in
	Brassica napus, Brassica rapa, and 80-99% homologues with the
	sequence in Gossypium raimondii, Populus trichocarpa, Cucumis


PhytoAB Inc.



sativus, Spinacia oleracea, Solanum tuberosum, Solanum lycopersicum, Zea mays, Panicum virgatum, Setaria viridis, Sorghum bicolor, Oryza sativa Japonica Group, Triticum aestivum, Hordeum vulgare.

For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 15 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4°C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only