

Anti-tRNA (cytosine(38)-C(5))-methyltransferase 2 antibody

Catalog: PHY3500A

Product Information

Description:	Rabbit polyclonal antibody
Background:	DNMT2 is a DNA methyltransferase homolog. Human Dnmt2 methylates
	tRNA-Asp and can methylate Arabidopsis tRNA-Asp in vitro.
Synonyms:	DNMT2, ATDNMT2, DNA METHYLTRANSFERASE 2, TRDMT1, DNA
	METHYLTRANSFERASE-2
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from
	Arabidopsis thaliana DNMT2 (AT5G25480).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb{C}$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at $4^\circ C$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	44 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immuniza <mark>tion is</mark> 100% homologues with the sequence in <i>Vitis</i>
	<i>vinifera, Brassica napus, Brassica rapa,</i> and 80-99% homologues
	with the sequence in Cucumis sativus, Gossypium raimondii, Populus


PhytoAB Inc.



trichocarpa, Oryza sativa, Hordeum vulgare, Triticum aestivum, Setaria viridis, Sorghum bicolor, Panicum virgatum, Zea mays, Glycine max, Medicago truncatula, Solanum lycopersicum, Solanum tuberosum, Nicotiana tabacum, Spinacia oleracea, Physcomitrium patens.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 15 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only