

## Anti-S-adenosyl-L-methionine-dependent methyltransferases superfamily protein antibody

Catalog: PHY3498A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** TRM4B is a cytosine-5--methyltransferase. Mutants have decreased methyl

cytosine and defects in root development.

**Synonyms:** TRM4B, ATTRM4B

**Immunogen:** KLH-conjugated synthetic peptide (15 aa from Central section) derived from

Arabidopsis thaliana TRM4B (AT2G22400).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70  $^{\circ}$ C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.


**Expected / apparent MW:** 90 kDa

**Predicted Reactivity:** For more species homologues information, please contact tech

support at tech@phytoab.com.



## **Application Example**



Recom: 2.5 ng, 15 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.